## 速報

## 干潟の海草藻場における ジュゴンの摂餌行動モニタリング

堤 千華\*<sup>1</sup> 市川光太郎\*<sup>1</sup> 赤松友成\*<sup>2</sup> 荒井修亮\*<sup>1</sup> 新家富雄\*<sup>3</sup> 原 武史\*<sup>4</sup> Kanjana Adulyanukosol\*<sup>5</sup>

- 1 京都大学大学院情報学研究科,〒606-8501 京都市左京区吉田本町
- ・2 独立行政法人水産総合研究センター水産工学研究所、〒 314-0408 茨城県神栖市波崎 7620-7
- ・3 株式会社システムインテック研究センター、〒 424-8610 静岡市清水区折戸3-20-1 東海大学フロンティアリサーチセンター内
- 4 社団法人日本水産資源保護協会。〒104-0054 東京都中央区勝どき2-18-1 黎明スカイレジテルビル西館303-2
- 5 Phuket Marine Biological Center P.O.BOX 60, Phuket 83000, Thailand

2005年7月4日受付, 2005年7月29日採録

## Abstract

Dugongs (Dugong dugon) are herbivorous marine mammals and endangered species. Unfortunately, the lack of basic research especially on use of seagrass patches by dugongs prevents us from taking any effective countermeasures for the conservation. In this study, to monitor feeding behavior of dugongs in the seagrass bed of a tidal flat, we propose a visual-acoustic combined observation method which is comprised of the visual observation and the acoustic observation using a digital camera and automatic underwater sound monitoring systems for dugong (AUSOMS-D). This new observation method was tested in Thai waters. The increase in the number of dugong feeding trails was observed during nighttime by the visual observation. Dugong calls and feeding sounds were recorded by the acoustic observation. The increase of the dugong feeding trails coincided with the feeding sounds. Therefore, the visual-acoustic combined observation method is effective in monitoring feeding behavior of dugongs. This method has technical benefits not to impact both on the dugongs and the seagrass beds.