

沿岸生態リスク評価モデルの開発

－瀬戸内海モデル－

江里口知己*¹ 市川哲也*¹ 桃井幹夫*¹ 酒井亨*¹ 中田喜三郎*² 堀口文男*³

- 1 株式会社 中電シーティーアイ, 環境情報部 〒450-0003 愛知県名古屋市中村区名駅南1-27-2
- 2 東海大学海洋学部, 〒424-8610 静岡市清水区折戸3-20-1
- 3 独立行政法人 産業技術総合研究所, 化学物質リスク管理研究センター 〒305-8569 茨城県つくば市小野川16-1

2006年6月15日受付, 2006年9月8日採録

Abstract

We have already developed ecological risk assessment models for Tokyo Bay and Ise Bay, called AIST-RAMTB and AIST-RAMIB. Using these models, users can easily estimate environmental concentrations of chemicals and their risk to marine organisms on Windows-based PCs. Based on these models, we developed a similar model suitable for the Seto-Inland Sea, called AIST-RAMSIS (National Institute of Advanced Industrial Science and Technology - The Risk Assessment Model for Seto-Inland Sea : AIST-RAMSIS). Environmental concentrations of chemicals can be estimated only by setting parameters of loading fluxes and calculation period using dialog-based graphic user interfaces (GUI). Using the estimated environmental concentrations of chemicals and their no observed effect concentrations (NOEC) for a target species, either risk to the species or an ecological risk can be determined. The results of environmental concentrations of chemicals and associated risk can be stored as CSV or image format files, and these results can be efficiently used or modified in other applications for other purposes.